



Deterministic OR Models

CO 370



Bertrand Guenin

Preface

Disclaimer Much of the information on this set of notes is transcribed directly/indirectly from the lectures of CO 370 during Fall 2021 as well as other related resources. I do not make any warranties about the completeness, reliability and accuracy of this set of notes. Use at your own risk.

What is operations research (OR)? There's no standard definitions for it. One particular definition: use of mathematical models to make complex decisions for real life problems. The origin is British military in WW2. OR is actually everywhere today. Key milestone: Simplex algorithm (1947).

Recall optimization problem is of the form:

$$\begin{array}{ll} \max & f(x) \\ \text{s.t.} & \text{a set of constraints} \end{array}$$

There are some applications: mail delivery, machine scheduling, inventory problem, network design, facility location, class scheduling, portfolio optimization, surgery planning, sensor location.

For any questions, send me an email via <https://notes.sibeliusp.com/contact>.

You can find my notes for other courses on <https://notes.sibeliusp.com/>.

Sibelius Peng

Contents

Preface	1
I Formulations	4
1 LP formulations	5
1.1 Production problem	5
1.2 Minimax	6
2 Flows	8
2.1 Max st-flow model	8
2.2 Min cost flow model	9
3 IP formulations	11
3.1 IP tricks	11
3.2 Modeling piece-wise linear function	11
3.3 Modeling union of polyhedra	12
3.4 Perfect formulations	12
3.5 Application to flows	15
4 Cone programming	17
4.1 Examples of cones	17
4.2 Cone programming model	18
4.3 (General) SOC program	18
5 Robust optimization	19
5.1 Model uncertainty for a	19
5.2 Modeling uncertainty by a ball	19
5.3 Modeling uncertainty by a hypercube	20
II Interpretations of optimal solutions	22
6 Duality review	23
6.1 Weak duality	23
6.2 Strong duality	23
6.3 Complementary Slackness conditions	23
7 Economic interpretation of dual variables	25
8 Sensitivity Analysis	27

8.1	Primal and dual Optimal basis	27
8.2	Review of simplex	28
8.3	Changes to the right-hand-side	29
8.4	Changes in the objective function	30
9	Primal and dual Simplex	32
9.1	Tableaus	33
9.2	Primal and Dual Simplex via tableaus	33
9.3	Parametric Linear Programs	34
III	Solving Optimization Techniques	36
10	Cutting planes - review	37
10.1	Cutting plane algorithm	37
11	Column generation	39
11.1	Column generation algorithm	39
11.2	Cutting stock	40



PART I:

FORMULATIONS

Picture from <https://www.catalent.com/oral-dose/development-bioavailability/formulation/>

LP formulations

1.1 Production problem

Products $J = \{1, \dots, n\}$

Resources $I = \{1, \dots, m\}$

Data:

- $\forall j \in J : c_j =$ value of unit of product j
- $\forall i \in I : b_i =$ number of units of resource i available
- $\forall i \in I, \forall j \in J : a_{ij} =$ number of units of resource i going to product j

Goal: maximize values of product made subject to available resources

Var: $x_j =$ number of units of product j produced

Then problem is

$$\begin{aligned} \max \quad & \sum_{j \in J} c_j x_j \\ \text{s.t.} \quad & \sum_{j \in J} a_{ij} x_j \leq b_i \quad (i \in I) \\ & x_j \geq 0 \quad (j \in J) \end{aligned}$$

Now let's generalize this problem to have more than one period.

Products $J = \{1, \dots, n\}$

Resources $I = \{1, \dots, m\}$

Periods $K = \{1, \dots, p\}$

Then we have **data**

- $\forall j \in J, k \in K : c_{jk} =$ unit value of product j in period k
- $\forall i \in I, k \in K : w_{ik} =$ unit price for resource i in period k
- $\forall i \in I, j \in J : a_{ij} =$ number of units of resource i going into product j

and the **goal:** decide how much of each resource to buy & how much of each product to make during each period, to maximize total profit. Unused resources are available at next time period.

Var:

- p_{ik} = number of units of resource i , purchased at start of period k
- x_{jk} = number of units of product j made in period k
- z_{ik} = number of units of resource i at the end of period k

$$\text{Profit} = \sum_{k \in K} \left[\sum_{j \in J} c_{jk} x_{jk} - \sum_{i \in I} w_{ik} p_{ik} \right] \tag{1.1}$$

Then we keep track of resources: for $i \in I, k \in K$

$$z_{ik} = z_{i(k-1)} + p_{ik} - \sum_{j \in J} a_{ij} x_{jk} \tag{1.2}$$

and we define for $i \in I$,

$$z_{i0} = 0 \tag{1.3}$$

Thus the optimization problem is

$$\begin{aligned} \max \quad & \text{(1.1)} \\ \text{s.t.} \quad & \text{(1.2), (1.3)} \\ & p, x, z \geq 0 \end{aligned} \tag{P}$$

Remark:

If (P) has a feasible solution of value that is bigger than 0, then (P) is unbounded. So we are missing some assumptions, maybe? For example, b_{ik} = amount of resource i that can be bound during period k . Then we can add constraints: $p_{ik} \leq b_{ik}$.

1.2 Minimax


Consider the problem of the form

$$\begin{aligned} \min_x \max \{ f_1(x), \dots, f_k(x) \} &:= g(x) \\ \text{s.t.} \quad & \dots \end{aligned}$$

where $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$.

Example:

$f_1(x) = -x - 1, f_2(x) = 0, f_3(x) = x - 1$. Then $\max\{f_1(x), f_2(x), f_3(x)\}$ is as follows



A motivation

- $\forall i \in [k], f_i(x) =$ completion time for task i .
- Project consists of task $1, \dots, k$.
- $g(x) =$ completion time of entire project

Note that minimax is not an optimization problem as we defined it. We can revise it as follows

$$\begin{array}{ll} \min & y \\ \text{s.t.} & y \geq f_1(x) \\ & y \geq f_2(x) \\ & \vdots \\ & y \geq f_k(x) \\ & \dots \end{array}$$

An application minimize a piece-wise linear convex function using linear programming.

Flows

digraph

A directed graph (digraph) is a pair (V, E) where

- V is a set of vertices,
- E is a set of ordered pairs of vertices called arcs.

Notation Let $q \in V$, then

$$\delta^+(q) = \{e \in E \mid e \text{ leaves } q\}$$

$$\delta^-(q) = \{e \in E \mid e \text{ arrives at } q\}$$

2.1 Max st-flow model

Given

1. digraph $G = (V, E)$,
2. two vertices $s, t \in V$, and $s \neq t$,
3. $\forall e \in E$, arc e has capacity $u_e \geq 0$.

Now we construct an LP.

For every arc e , we will have a variable x_e , and x_e will be called the **flow** on arc e .

Notation Let $q \in V$:

$$f_x(q) := \sum_{e \in \delta^+(q)} x_e - \sum_{e \in \delta^-(q)} x_e$$

The maximization problem is then

$$\begin{aligned} \max \quad & f_x(s) \\ \text{s.t.} \quad & f_x(q) = 0 \quad (q \in V, q \neq s, q \neq t) \\ & 0 \leq x_e \leq u_e \quad (e \in E) \end{aligned} \tag{P}$$

A feasible solution to (P) is a flow. An optimal solution to (P) is a maximum flow. The value of a flow x is $f_x(S)$.

Remark:

(P) is always feasible. It is not unbounded, so there always exists a maximum flow.

Application computer network. Suppose we have

- computers s and t ($s \neq t$)
- capacity u_e (gb/s) for every link e

The goal is to computer number gb that can be sent from s to t across network. Then x_e is the amount of information across e . $f_x(q) = 0$ means no information lost.

Magic property

If u is integer, then there exists an optimal solution to (P) that is integer.

Remark:

We need the condition “ u is integer” so that the property is still true. Also, an optimal solution to (P) is not necessarily integers.

Generalize max st-flows

We can add lower bounds to arcs: $\ell_e \forall e \in E$

$$\begin{aligned} \max \quad & f_x(s) \\ \text{s.t.} \quad & f_x(q) = 0 \quad (q \in V, q \neq s, q \neq t) \\ & \ell_e \leq x_e \leq u_e \quad (e \in E) \end{aligned} \tag{Q}$$

Magic property - revised

If ℓ, u is integer, and there exists an optimal solution to (Q), then there exists an optimal solution to (Q) that is integer.

Example: Consistent rounding

The goal is to round all entries to nearest up/down integer, so that row sums & column sums still hold.

Any feasible solutions give consistent rounding.

2.2 Min cost flow model

Given

- Digraph $G = (V, E)$
- Capacities $u_e \geq 0$ ($e \in E$)
- Costs c_e ($e \in E$)
- Supply/demands b_q ($q \in V$)

Then the model is

$$\begin{array}{ll} \min & \sum_{e \in E} c_e x_e \\ \text{s.t.} & f_x(q) = b_q \quad (q \in V) \\ & 0 \leq x_e \leq u_e \quad (e \in E) \end{array} \quad (\text{P})$$

Similarly, feasible solution to (P) is a flow. An optimal solution to (P) is a min cost flow.

Magic property - min cost flows

Suppose u, b are integer, then if there exists a min cost flow, then there exists a min cost flow that is integer.

Similarly, we can add a lower bound:

$$\begin{array}{ll} \min & \sum_{e \in E} c_e x_e \\ \text{s.t.} & f_x(q) = b_q \quad (q \in V) \\ & \ell_e \leq x_e \leq u_e \quad (e \in E) \end{array} \quad (\text{P})$$

Magic property - min cost flows (revised)

Suppose u, b, ℓ are integer, then if there exists a min cost flow, then there exists a min cost flow that is integer.

What is a necessary condition for b_q so that there exists a flow? $\sum_{q \in V} b_q = 0$.

Example: Staffing problem

hours	# of employees required
1-2	2
2-3	3
3-4	3
4-5	2

and we have cost of hiring a single employee between

hours	cost
1-5	6
1-4	4
3-5	5
2-4	3

The goal is to minimize cost of hiring employees while meeting staff needs.

3

IP formulations

3.1 IP tricks

Imaging we are forcing variable to take some prescribed set of values: $x \in \{5, 9, 13, 36\}$. Then we can introduce variables $z_1, z_2, z_3, z_4 \in \{0, 1\}$ so that we have two constraints:

$$\begin{aligned}1 &= z_1 + z_2 + z_3 + z_4 \\x &= 5z_1 + 9z_2 + 13z_3 + 36z_4\end{aligned}$$

3.2 Modeling piece-wise linear function

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise linear. Given a_1, \dots, a_k and f_1, \dots, f_k such that $f(a_i) = f_i$. The goal is to write IP constraints with variables x, y such that $y = f(x)$, $x \in [a_1, a_k]$.

To generalize,

\uparrow
dom(f)

1. $\lambda_1, \dots, \lambda_k \geq 0, \sum_{i=1}^k \lambda_i = 1$
2. $x = \sum_{i=1}^k \lambda_i a_i$
3. $y = \sum_{i=1}^k \lambda_i f_i$
4. $z_0, \dots, z_k \in \{0, 1\}, z_0 = z_k = 0$
5. $\sum_{i=0}^k z_i = 1$
6. $\forall p \in [k], \lambda_p \leq z_{p-1} + z_p$

By (4) and (5), we may assume that $z_p = 1$ and $z_j = 0 \forall j \neq p$.

Claim λ_p, λ_{p+1} are the only non-zero λ variables.

Proof:

Pick $j \neq p, j \neq p+1$. As

$$0 \leq \lambda_j \leq z_{j-1} + z_j = 0 + 0 = 0$$

□

With this claim, we can simplify (2) and (3).

3.3 Modeling union of polyhedra

Let

$$P_1 = \{x \mid A^1x \leq b^1\}$$

$$P_2 = \{x \mid A^2x \leq b^2\}$$

Goal: write condition: $x \in P_1 \cup P_2$ as part of IP.

Hypothesis: If $x \in P_1 \cup P_2$, then $0 \leq x \leq U$ for some U .

Constraints:

1. $y_1, y_2 \in \{0, 1\}$
2. $y_1 + y_2 = 1$
3. $x = x^1 + x^2$
4. $A^i x^i \leq y_i b^i, i = 1, 2$
5. $0 \leq x^i \leq y_i U, i = 1, 2$

To show that $x \in P_1 \cup P_2 \iff \exists x^1, x^2, y_1, y_2$ such that all 5 conditions hold.

Proof:

First let's assume (1) - (5) hold. We may assume $y_1 = 1, y_2 = 0$. Then (5) tells us $x^2 = 0$. (3) tells us $x = x^1$. (4) implies

$$A^1 x^1 \leq y_1 b^1 \implies A^1 x \leq b^1 \implies x \in P_1 \subseteq P_1 \cup P_2$$

Now suppose $x \in P_1 \cup P_2$. We may assume $x \in P_1$. Then we set $y_1 = 1, y_2 = 0, x^1 = x, x^2 = 0$. Then we can verify that all 5 conditions hold. \square

3.4 Perfect formulations

basis

Let A be a matrix with column indices $\{1, \dots, n\}$. Then $B \subseteq \{1, \dots, n\}$ is a basis if

1. A_B square,
2. A_B non-singular.

Remark:

A has a basis \iff rows of A are independent.

basic solution

Let A be a matrix with column indices $1, \dots, n$. Consider

$$Ax = b \tag{*}$$

Pick B as basis of A . Then x is a basic solution of (*) if

1. $Ax = b$.
2. $x_j = 0$ if $j \notin B$.

x is a basic solution for (*) if x is a basic solution for some basis B .

standard equality form

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

and rows of A are independent.

The correctness of simplex algorithm implies the following theorem:

Theorem 3.1

If an LP is SEF has an optimal solution, then it has an optimal solution that is basic.

Let A be matrix, b vector with same number of entries as rows of A . Then $A \leftarrow_j b$ denotes matrix obtained from A by replacing column j by b .

Theorem 3.2: Cramer's rule

Let M be a non-singular matrix and consider $Mx = b$.

$$\bar{x}_j = \frac{\det(M \leftarrow_j b)}{\det(M)} \quad \forall j$$

then $M\bar{x} = b$.

Proposition 3.3

Let M be square matrix with $\det(M) = \pm 1$, and M, b are integer. Then there exists a unique solution to $Mx = b$ is integer.

Proof:

Directly from Cramer's rule. □

Totally unimodular matrix

A matrix A is totally unimodular if every square submatrix N of A , $\det(N) \in \{0, +1, -1\}$.

Proposition 3.4

Let A be TU, b integer. Then every basic solution of $Ax = b$ is integer.

Proof:

Suppose \bar{x} is a basic solution for basis B . If $j \notin B$, $\bar{x}_j = 0$, which is integer. The basic variables \bar{x}_B are the unique solution to $A_B \bar{x}_B = b$. Since A is TU, then $\det(A_B) \in \{0, -1, +1\}$. Since B is basis, $\det(A_B) \neq 0$. Then $\det(A_B) = \pm 1$. A_B is integer as it is TU. Then by Proposition 3.3, \bar{x}_B is integer. □

Theorem 3.5

If an LP is SEF has an optimal solution, it has an optimal solution that is basic.

Theorem 3.6: main theorem

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b, x \geq 0 \end{aligned} \quad (\text{P})$$

Suppose A is TU and b integer. Then if (P) has an optimal solution, it has an optimal solution that is integer.

Proof:

We may assume rows of A are linearly independent. Then (P) is SEF. Thus if (P) has an optimal solution, then by Theorem 3.5, it has an optimal basic solution \bar{x} . By Proposition 3.4, every basic solution is integer. In particular, so is \bar{x} . \square

Constructing TU matrices**Proposition 3.7**

Let A be a $0, -1, +1$ matrix such that for every column

1. we have at most one $+1$,
2. we have at most one -1 .

Then A is TU.

Proof:

We need to show every $k \times k$ submatrix M of A has $\det(M) \in \{0, -1, +1\}$. We can prove it by induction on k . Discuss by cases.

- M has a column of zeros. Then $\det(M) = 0$.
- Every column has exactly one $+1$ and one -1 , then row sum is zero, then rows are linearly dependent, thus $\det(M) = 0$.
- One column has a unique non-zero entry M_{ij} . Applying cofactor expansion,

$$\det(M) = M_{ij}(-1)^{i+j} \det(M'_{ij})$$

Then by induction, the statement holds. \square

Constructing TU matrices from TU matrices

Proposition 3.8

Let A be TU. Then

1. permutation of A ,
 2. A^T ,
 3. any matrix obtained from A by multiplying rows/columns by -1 ,
 4. any matrix obtained by adding a unit vector to the rows/columns
- are TU.

Theorem 3.9

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b, x \geq 0 \end{aligned} \tag{P}$$

Suppose A is TU, b integer. Then if (P) has an optimal solution, then it has one that is integer.

Theorem 3.10

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b, 0 \leq x \leq U \end{aligned} \tag{Q}$$

Suppose A is TU, b and U are integer. Then if (Q) has optimal solution, then it has one that is integer.

Proof:

We can rewrite (Q) in SEF:

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & \begin{bmatrix} A & 0 \\ I & I \end{bmatrix} \begin{bmatrix} x \\ s \end{bmatrix} = \begin{bmatrix} b \\ U \end{bmatrix} \\ & x, s \geq 0 \end{aligned} \tag{P'}$$

We denote the first constraint by $A'x' = b'$. Now note that b' is integer as b, U are integer. It's clear that A' is TU. Then apply Theorem 3.9. \square

3.5 Application to flows

vertex-arc incidence matrix

M is the vertex-arc incidence matrix of digraph $G = (V, E)$ if

1. rows of M correspond to V ,
2. columns of M correspond to E ,
3. for column uv we have $+1$ for every u , -1 for entry v and 0 otherwise.

Remark:

Let M be vertex-arc incidence matrix of $G = (V, E)$. Pick $v \in V$: for every entry e of row v , if $e \in \delta^+(v)$, then we have $+1$; if $e \in \delta^-(v)$, we have -1 , otherwise we have 0 .

Magic property - min cost flow

$$\begin{aligned} \min \quad & c^T x \\ \text{s.t.} \quad & f_x(q) = b_q \quad (q \in V) \\ & 0 \leq x \leq U \end{aligned} \tag{P}$$

Suppose b, U integer, and there exists an optimal solution. Then there exists an integer optimal solution.

Proof:

Let A be the vertex-arc matrix.

Claim $f_x(q) = b_q \iff \text{row}_q(A)x = b_q$.

Proof of the claim:

Using the remark, we know that

$$\text{row}_q(A)x = \sum_{e \in \delta^+(q)} x_e - \sum_{e \in \delta^-(q)} x_e := f_x(q)$$

□

We can write (P) as

$$\begin{aligned} \min \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & 0 \leq x \leq U \end{aligned}$$

We proved A is TU. Then it follows by Theorem 3.10.

□

4

Cone programming

This is generalization of

1. linear programming
2. second-order cone programming
3. semi-definite programming

We have good algorithms for these problems.

cone

$C \subseteq \mathbb{R}^m$ is a cone if

1. C is convex,
2. $\forall \lambda \geq 0$ and $a \in C, \lambda a \in C$.

The second condition: C is closed under (non-negative scaling).

pointed cone

A cone is pointed if it does not contain an infinite line.

4.1 Examples of cones

$$\mathbb{R}_+^m = \{u \in \mathbb{R}^m : u \geq 0\}$$

Second Order Cone (SOC):

$$L^m = \{u \in \mathbb{R}^m : \|(u_1, \dots, u_{m-1})\| \leq u_m\}$$

semi-definite matrix

An $n \times n$ matrix M is semi-definite if $\forall x \in \mathbb{R}^n: x^T M x \geq 0$.

Semi-definite cone is

$$S_+^n = \{n \times n \text{ semi-definite matrices } M\}$$

Direct sum

Let $A \subseteq \mathbb{R}^p, B \subseteq \mathbb{R}^q$. The direct sum of A and B is

$$A \oplus B := \{(a, b) : a \in A, b \in B\} \subseteq \mathbb{R}^{p+q}$$

Proposition 4.1

Let A, B be cones, then $A \oplus B$ is a cone.

4.2 Cone programming model

Let $c \in \mathbb{R}^n, A$ $m \times n$ matrix, $b \in \mathbb{R}^m, K$ pointed cone.

$$\begin{aligned} \min \quad & c^T x \\ \text{s.t.} \quad & Ax - b \in K \end{aligned} \tag{P}$$

Cases:

1. $K = \mathbb{R}_+^m$. Then $Ax - b \in \mathbb{R}_+^m \iff Ax \geq b$. Then it's an LP.
2. $K = L^m$. $Ax - b \in L^m$ can be rewritten as

$$\begin{aligned} \begin{bmatrix} A' \\ d^T \end{bmatrix} x - \begin{bmatrix} b' \\ p \end{bmatrix} \in L^m \\ \|A'x - b'\| \leq d^T x - p \end{aligned}$$

and this is restricted SOC program.

3. $K = S_+^n$. This is semi-definite program.

Example: Least square problem

$$\begin{aligned} \min_x \quad & \|Ax - b\| \\ \text{or} \quad & \\ \min \quad & t \\ \text{s.t.} \quad & \|Ax - b\| \leq t \end{aligned}$$

which is restricted SOC program

4.3 (General) SOC program

$$\begin{aligned} \min \quad & c^T x \\ \text{s.t.} \quad & \|A_i x - b_i\| \leq d_i^T x - p_i \quad (i = 1, \dots, k) \end{aligned}$$

Is this still a cone program? In other words, is there A, B, K such that the above constraint is equivalent to $Ax - b \in K$? Yes. It's direct sum of pointed cones.

5

Robust optimization

$$\begin{array}{ll} \min & \dots \\ \text{s.t.} & a^T x \leq \beta \\ & \dots \end{array}$$

This particular constraint might be important. And a is part of data that is given, but a here might have some imprecision. This brings us the uncertainty.

5.1 Model uncertainty for a

Given: $\bar{a} \in \mathbb{R}^n$ estimate, $\epsilon \subseteq \mathbb{R}^n$ uncertainty. True a satisfies $a - \bar{a} \in \epsilon$.

Then we can write the condition

$$a^T x \leq \beta \quad (a - \bar{a} \in \epsilon)$$

We have different choices for ϵ , or think it for different ways.

For example, the ball:

$$a^T x \leq \beta \quad (a - \bar{a} \in \{u : \|u\| \leq r\} := \epsilon_1) \tag{5.1}$$

the cube:

$$a^T x \leq \beta \quad (a - \bar{a} \in \{u : -r \leq u_i \leq r, \forall i\} := \epsilon_2) \tag{5.2}$$

Let

$$S_1 = \{x \mid x \text{ satisfies (5.1)}\}$$

$$S_2 = \{x \mid x \text{ satisfies (5.2)}\}$$

What's the relationship between S_1 and S_2 ? We know that $\epsilon_1 \subseteq \epsilon_2$. The constraints in (5.1) form a subset of constraints in (5.2). Thus $S_2 \subseteq S_1$.

We will see: (5.1) can be replaced by a single SOC constraint: $\|f(x)\| \leq g(x)$ where f, g affine. (5.2) can be replaced by a finite number of linear constraints (with additional variables).

5.2 Modeling uncertainty by a ball

Given data $r > 0$, $\bar{a} \in \mathbb{R}^n$, $\beta \in \mathbb{R}$. We then have $\epsilon = \{u \in \mathbb{R}^n \mid \|u\| \leq r\}$.

$$a^T x \leq \beta \quad (a - \bar{a} \in \epsilon) \tag{*}$$

As (*) has infinitely many of inequalities, the goal is to replace (*) with a single inequality. The key idea is to fix x . Note that

$$\begin{aligned} (*) &\iff \max\{a^T x : a - \bar{a} \in \epsilon\} \leq \beta \\ &\bar{a}^T x + \max\{a^T x - \bar{a}^T x : a - \bar{a} \in \epsilon\} \leq \beta \\ &\bar{a}^T x + \max\{(a - \bar{a})^T x : a - \bar{a} \in \epsilon\} \leq \beta \\ &\bar{a}^T x + \max\{u^T x : u \in \epsilon\} \leq \beta \end{aligned}$$

Note that $\max\{u^T x : u \in \epsilon\} = r\|x\|$ given ϵ is a radius r ball. Then

$$\bar{a}^T x + r\|x\| \leq \beta$$

and this is exactly SOC constraint.

ellipsoid

An ellipsoid is a set of the form $\{Pu : \|u\| \leq 1\}$ and P is a non-singular matrix.

For example, let $P = \begin{bmatrix} 2 & 0 \\ 0 & 1 \end{bmatrix}$

This is generalization of the ball.

5.3 Modeling uncertainty by a hypercube

Recall the constraint

$$a^T x \leq \beta \quad (a - \bar{a} \in \epsilon) \quad (\heartsuit)$$

where $\epsilon = \{U \mid -r \leq v_j \leq r, \forall j\}$. We need to show

1. (\heartsuit) is equivalent to single constraint, involving absolute values.
2. (\heartsuit) is equivalent to finite number of linear constraints.

$$a^T x \geq \beta \quad (a - \bar{a} \in \epsilon) \quad (\heartsuit)$$

Fix x . Then the above constraint is equivalent to

$$\min\{a^T x \mid a - \bar{a} \in \epsilon\} \geq \beta$$

Then similarly,

$$\bar{a}^T x + \min\{u^T x \mid u \in \epsilon\} \geq \beta$$

Note that assume u is optimal, then

$$\begin{aligned} u^T x &= \sum_j u_j x_j \\ &= \sum_{j:x_j \geq 0} u_j x_j + \sum_{j:x_j < 0} u_j x_j \\ &= \sum_{j:x_j \geq 0} -r x_j + \sum_{j:x_j < 0} r x_j \\ &= -r \sum_j |x_j| \end{aligned}$$

Proposition 5.1

$$a^T x \geq \beta \quad (a - \bar{a} \in \{u \mid -r \leq u_j \leq r, \forall j\})$$

is equivalent to

$$\bar{a}^T x - r \sum_j |x_j| \geq \beta$$

Corollary 5.2

$$a^T x \leq \beta \quad (a - \bar{a} \in \{u \mid -r \leq u_j \leq r, \forall j\})$$

is equivalent to

$$\bar{a}^T x + r \sum_j |x_j| \leq \beta$$

Proposition 5.3

1. $\exists x$ such that $\bar{a}^T x + r \sum_j |x_j| \leq \beta$
2. $\exists x, y$ such that $\bar{a}^T x + r \sum_j y_j \leq \beta$, $y_j \geq x_j$, $y_j \geq -x_j$.

are equivalent.

Proposition 5.4

$$a^T x \leq \beta \quad (a - \bar{a} \in \{u \mid -r \leq u_j \leq r, \forall j\})$$


is equivalent to

$$\begin{aligned} \bar{a}^T x + r \sum_j y_j &\leq \beta \\ y_j &\geq x_j, y_j \geq -x_j, \forall j \end{aligned}$$

This can be generalized to be a rectangle. Given $\Delta \geq 0$. Then

$$\epsilon = \{i \mid -\delta_j \leq u_j \leq \delta_j, \forall j\}$$

For example, $\Delta = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$



PART II:

INTERPRETATIONS OF OPTIMAL SOLUTIONS

Picture from <https://leadingincontext.com/2019/11/06/what-is-duality/>

Duality review

$$\begin{array}{ll}
 \max & c^T x \\
 \text{s.t.} & Ax \leq b \\
 & x \geq 0
 \end{array} \quad (\text{P}_{\max})$$

$$\begin{array}{ll}
 \min & b^T y \\
 \text{s.t.} & A^T y \leq c \\
 & y \geq 0
 \end{array} \quad (\text{P}_{\min})$$

Then we have this table

P_{\max}	P_{\min}
$x_j \geq 0$	constraint $j \geq$
$x_j \leq 0$	constraint $j \leq$
x_j free	constraint $j =$
constraint $j \leq$	$y_i \geq 0$
constraint $j \geq$	$y_i \leq 0$
constraint $j =$	y_i free

6.1 Weak duality

Let P_{\max} and P_{\min} be a primal/dual pair. Let \bar{x} and \bar{y} feasible for P_{\max} and P_{\min} respectively, then $c^T \bar{x} \leq b^T \bar{y}$.

Then if $c^T \bar{x} = b^T \bar{y}$, then \bar{x} optimal for P_{\max} , \bar{y} optimal for P_{\min} .

6.2 Strong duality

Let P_{\max} and P_{\min} be a primal/dual pair.

- P_{\max} has an optimal solution \bar{x} iff P_{\min} has an optimal solution \bar{y} .
- If (1) holds, then $c^T \bar{x} = b^T \bar{y}$.

Strong duality is what makes weak duality useful.

6.3 Complementary Slackness conditions

\bar{x}, \bar{y} satisfy CS conditions if

1. $\forall j: \bar{x}_j = 0$ or constraint j of P_{\min} tight.
2. $\forall i: \bar{y}_i = 0$ or constraint i of P_{\max} tight.

CS theorem

Let \bar{x}, \bar{y} feasible for P_{\max} and P_{\min} . Then the following are equivalent:

1. \bar{x}, \bar{y} optimal for P_{\max} and P_{\min} .
2. CS conditions hold.
3. $c^T \bar{x} = b^T \bar{y}$.

Economic interpretation of dual variables

Recall the production problem

1. Products $J = \{1, \dots, n\}$, Resources $I = \{1, \dots, m\}$.
2. $\forall j \in J: c_j =$ unit value of product j .
3. $\forall i \in I: b_i =$ number of units of resource i available.
4. $\forall i \in I, j \in J: A_{ij} =$ number of units of resource i in one unit of product j .

The formulation is

$$\begin{aligned} \max \quad & \sum_{j \in J} c_j x_j \\ \text{s.t.} \quad & \sum_{j \in J} A_{ij} x_j \leq b_i \quad (i \in I) \\ & x_j \geq 0 \quad (j \in J) \end{aligned}$$

Now we can write it in a more compact way:

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax \leq b \\ & x \geq 0 \end{aligned} \tag{P}$$

and its dual

$$\begin{aligned} \min \quad & b^T y \\ \text{s.t.} \quad & A^T y \geq c \\ & y \geq 0 \end{aligned} \tag{D}$$

What is (D) doing? Consider a special case.

Let x_1 be number of hammer, x_2 be number of pliers. The formulation is

$$\begin{aligned} \max \quad & \begin{pmatrix} 130 & 100 \end{pmatrix} x \\ \text{s.t.} \quad & \begin{pmatrix} 1.5 & 1 \\ 1 & 1 \\ 0.3 & 0.5 \end{pmatrix} x \leq \begin{pmatrix} 27 \\ 21 \\ 9 \end{pmatrix} \end{aligned}$$

where three rows represent steel, plastic, time respectively. Primal optimal solution $\bar{x} = (12, 9)^T$. Its dual

$$\begin{aligned} \min \quad & \begin{pmatrix} 27 & 21 & 9 \end{pmatrix} y \\ \text{s.t.} \quad & \begin{pmatrix} 1.5 & 1 & 0.3 \\ 1 & 1 & 0.5 \end{pmatrix} y \geq \begin{pmatrix} 130 \\ 100 \end{pmatrix} \\ & y \geq 0 \end{aligned}$$

Then the dual solution $\bar{y} = (60 \ 40 \ 0)^T$. The dual variables attach to each of resources.

In general, we can think of $y_i =$ “price” of resource i . But this is not quite precise enough. More precisely, y_i is the lowest unit price at which resource i can be sold without losing money. This can also be called as market price, fair price, shadow price.

This is indeed the right interpretation. Let us look at constraint of (D):

- $y \geq 0$: prices are non negative
- $A^T y \geq c$:

$$\begin{aligned} \text{col}_j(A)^T y &\geq c_j \\ \sum_{i \in I} A_{ij} y_i &\geq c_j \end{aligned}$$

where c_j is unit value for product j . And the sum means the market value of resources going into one unit of product j .

Note that $b^T y = \sum_{i \in I} b_i y_i$ is the total market value of all resources. Then strong duality tells us total production value = market value of all resources.

Now let’s look at CS. Suppose \bar{x}, \bar{y} optimal for (P) and (D).

- $\forall i \in I, \text{row}_i(A)\bar{x} < b_i \implies \bar{y}_i = 0$. It tells us if we have excess resource i , then market value of resource i is zero.
- $\forall j \in J, \text{col}_j(A)^T \bar{y} > c_j \implies x_j = 0$. It tells us if market value of resource into one unit of product j is bigger than value of product j , then do not produce product j .

Proposition 7.1

Let $\alpha \geq 0$. Consider

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax \leq b \\ & x \geq 0 \end{aligned} \tag{P}$$

and

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax \leq b + \alpha e_i \\ & x \geq 0 \end{aligned} \tag{Q}$$

Suppose (P) has an optimal solution and let \bar{y} be the optimal solution to the dual of (P).

1. (Q) has an optimal solution.
2. $\text{OPT}(Q) \leq \text{OPT}(P) + \alpha \bar{y}_i$.

Proof:

Denote the duals of (P) and (Q) by (D) and (R).

Since (P) feasible and $\alpha \geq 0$, then (Q) feasible. \bar{y} feasible for (D), then \bar{y} feasible for (R). Then (Q) and (R) both have an optimal solution.

To prove (2), note that

$$\begin{aligned} \text{OPT}(Q) &\leq (b + \alpha e_i)^T \bar{y} && \text{weak duality} \\ &= b^T \bar{y} + \alpha \bar{y}_i \\ &= \text{OPT}(P) + \alpha \bar{y}_i && \text{strong duality} \end{aligned}$$

□

Sensitivity Analysis

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

Let B be an optimal basis. For what changes to A, b or c does B remain optimal?

8.1 Primal and dual Optimal basis

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

and dual

$$\begin{aligned} \max \quad & b^T y \\ \text{s.t.} \quad & A^T y \geq c \end{aligned} \tag{D}$$

Let B be basis of A , and N column indices of A not in B . Then this means we can partition $A = [B \mid N]$.

primal/dual basic solution

x is a primal basic solution if $x_B = A_B^{-1}b, x_N = 0$. y is a dual basic solution if $y = A_B^{-T}c_B$.

primal/dual feasible

B primal feasible if x feasible for (P). B dual feasible if y feasible for (D).

optimal basis

B is optimal if it is primal and dual feasible.

We now need to be convinced that this is a good definition. Now we can characterize primal/dual feasible basis.

Remark:

Let B be basis. Define $\bar{b} = A_B^{-1}b$. Then B is primal feasible if and only if $\bar{b} \geq 0$.

Remark:

Let B be basis. Define $y = A_B^{-T}c_B$, $\bar{c} = c - A^T y$. Then the following are equivalent:

- B dual feasible.
- $\bar{c} \leq 0$.
- $\bar{c}_N \leq 0$.

Proposition 8.1

If B is optimal basis then

1. primal basic solution is optimal,
2. dual basic solution is optimal.

Proof:

By CS theorem, it suffices to check CS conditions hold for x, y . □

Proposition 8.2

If an LP in SEF has an optimal solution, then it has an optimal basis.

8.2 Review of simplex

canonical form

(P) is in canonical form for some basis B if $A_B = I$ and $c_B = 0$.

Proposition 8.3

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

Let B be basis.

$$\begin{aligned} \max \quad & \bar{c}^T x + b^T y \\ \text{s.t.} \quad & A_B^{-1} Ax = \bar{b} \\ & x \geq 0 \end{aligned} \tag{Q}$$

where

$$\bar{b} = A_B^{-1}b, \quad \bar{c} = c - A^T y, \quad y = A_B^{-T}c_B$$

Then (Q) is in canonical form for B .

Simplex algorithm always STOP with an optimal basis.

8.3 Changes to the right-hand-side

Proposition 8.4

Let $\alpha \geq 0$. Consider

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

and

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b + \alpha e_i \\ & x \geq 0 \end{aligned} \tag{Q}$$

Suppose B optimal for (P). Then B optimal for (Q) if and only if

$$\bar{b} + \alpha \cdot \text{col}_i(A_B^{-1}) \geq 0$$

where $\bar{b} = A_B^{-1}b$.

The set of possible α here is the allowable range.

Proof:

Consider the duals

$$\begin{aligned} \min \quad & b^T y \\ \text{s.t.} \quad & A^T y \geq c \end{aligned} \tag{D_P}$$

$$\begin{aligned} \min \quad & (b + e_i)^T y \\ \text{s.t.} \quad & A^T y \geq c \end{aligned} \tag{D_Q}$$

B optimal for (P) means:

$$(1) \quad A_B^{-1}b \geq 0$$

$$(2) \quad \bar{c} = c - A^T y \leq 0, \quad y = A_B^{-T} c_B$$

B optimal for (Q) means:

$$(1') \quad A_B^{-1}(b + \alpha e_i) \geq 0$$

(2') same as (2) above.

We know (1) and (2) hold. When does (1') hold? Because (2') always hold. It holds when

$$A_B^{-1}(b + \alpha e_i) = A_B^{-1}b + \alpha A_B^{-1}e_i = \bar{b} + \alpha \text{col}_i(A_B^{-1}) \geq 0$$

□

Proposition 8.5

Let $\alpha \geq 0$. Consider

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

and

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b + \alpha e_i \\ & x \geq 0 \end{aligned} \tag{Q}$$

Suppose B optimal for (P). Let \bar{y} be the basic dual solution for (P) and basis B . Suppose α is in the allowable range. Then

$$\text{OPT}(Q) = \text{OPT}(P) + \alpha \bar{y}_i$$

Proof:

Consider duals. □

8.4 Changes in the objective function**Proposition 8.6**

Consider

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

and

$$\begin{aligned} \max \quad & d^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{Q}$$

where $d = c + \alpha e_j$ where $j \notin B$. Suppose B is optimal for (P). Then B optimal for (Q) iff $\alpha \leq -\bar{c}_j$.

Proof:

Similar as before. □

Proposition 8.7

Consider

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

and

$$\begin{aligned} \max \quad & d^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{Q}$$

where $d = c + \alpha e_j$ where $j \in B$. Suppose B is optimal for (P). Then B optimal for (Q) iff

$$\forall k \notin B : \bar{c}_k - \alpha \cdot \text{row}_r(A_B^{-1}) \text{col}_k(A) \leq 0.$$

And j corresponds to the r^{th} basis element.

For example, let $B = \{1, 4, 9, 12, 15\}$. Then for $j = 9$, we have $r = 3$.

Proposition 8.8

Consider

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax = b \\ & x \geq 0 \end{array} \quad (\text{P})$$

and

$$\begin{array}{ll} \max & d^T x \\ \text{s.t.} & Mx = b \\ & x \geq 0 \end{array} \quad (\text{Q})$$

where

$$M = \left[\begin{array}{c|c|c|c} A_1 & \cdots & A_n & f \end{array} \right], \quad d = (c_1, c_2, \dots, c_n, p).$$

Suppose B optimal for (P) and y basic dual solution for (P).. Then B optimal for (Q) iff $y^T f \geq p$.

Primal and dual Simplex

The idea is to solve dual problem. Consider two applications. One is **parametric linear programs**, which we can think of as extension of sensitivity analysis. Another is cutting planes.

(Primal) Simplex algorithm

At each step,

1. primal feasible basis B .
2. rewrite in canonical form for B .
3. if B dual feasible, then STOP.
4. if primal unbounded, then STOP.
5. find better basis: $B \cup \{j\} \setminus \{r\}$.

Dual Simplex algorithm

At each step,

1. dual feasible basis B .
2. rewrite in canonical form for B .
3. if B primal feasible, then STOP.
4. if dual unbounded, then STOP.
5. find better basis: $B \cup \{j\} \setminus \{r\}$.

Key ingredients:

- decide what variable enters
- decide what variable leaves
- detect unboundedness

Now we are going to show how to rewrite in canonical using notation of tableaus.

9.1 Tableaus

$$\begin{aligned} \max \quad & z = c^T x + \bar{x} \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned}$$

can be transformed into

$$\begin{aligned} z - c^T x &= \bar{z} \\ Ax &= b \end{aligned}$$

Then we have the tableau representation:

$$\begin{array}{c|cc} & z & x & \text{RHS} \\ \hline 1 & & -c^T & \bar{z} \\ \hline 0 & & & \\ \vdots & & A & b \\ \hline 0 & & & \end{array}$$

Remark:

Pivot on (i, j) . j enters, r leaves where r is the i^{th} basic variable.

9.2 Primal and Dual Simplex via tableaus

$$\begin{aligned} \max \quad & c^T x \\ \text{s.t.} \quad & Ax = b \\ & x \geq 0 \end{aligned} \tag{P}$$

$$\begin{aligned} \max \quad & b^T y \\ \text{s.t.} \quad & A^T y \geq c \end{aligned} \tag{D}$$

We have basis B . (P) rewritten in canonical form:

$$\begin{aligned} \max \quad & \bar{c}^T x + \bar{z} \\ \text{s.t.} \quad & \bar{A}x = \bar{b} \\ & x \geq 0 \end{aligned}$$

and put into tableau

$$\begin{array}{c|cc} 1 & -\bar{c}^T & \bar{z} \\ \hline 0 & & \\ \vdots & \bar{A} & \bar{b} \\ \hline 0 & & \end{array}$$

Primal Simplex B primal feasible.

1. if $\bar{c} \leq 0$, then STOP.
2. pick j such that $\bar{c}_j > 0$.
3. if $\text{col}_j(\bar{A}) \leq 0$, then STOP, (P) unbounded.
4. let i be the k index minimizing $\min_k \left\{ \frac{\bar{b}_k}{\bar{A}_{kj}} : \bar{A}_{kj} > 0 \right\}$
5. pivot on (i, j) .

Dual Simplex assume dual feasible, then $\bar{c} \leq 0$

1. if $\bar{b} \geq 0$, then STOP.
2. Pick i such that $\bar{b}_i < 0$
3. If $\text{row}_i(\bar{A}) \geq 0$, then STOP, (D) unbounded.
4. Let j be the k index minimizing $\min_k \left\{ \frac{\bar{c}_k}{\bar{A}_{ik}} : \bar{A}_{ik} < 0 \right\}$
5. Pivot on (i, j) .

Claim In (3) if $\text{row}_i(\bar{A}) \geq 0$, then (D) is unbounded.

Proof:

Note that

$$\underbrace{\bar{b}_i}_{<0} = \underbrace{\text{row}_i(\bar{A})}_{\geq 0} \underbrace{x}_{\geq 0}$$

Thus (P) infeasible. Then (D) is either infeasible or unbounded. But (D) is feasible. \square

Claim B remains dual feasible during each iteration.

Proof:

For the initial iteration, if we write out the tableau, $-\bar{c}^T \geq 0$, then dual feasible. For the next iteration, we want to show $-c' \geq 0$, thus dual feasible.

$$-c'_k = -\bar{c}_k + \frac{\bar{c}_j}{\bar{A}_{ij}} \bar{A}_{ik} \stackrel{?}{\geq} 0$$

and this is equivalent to check whether

$$\frac{\bar{c}_j}{\bar{A}_{ij}} \bar{A}_{ik} \geq \bar{c}_k$$

We then discuss by cases: $\bar{A}_{ik} \geq 0$ or $\bar{A}_{ij} < 0$. \square

9.3 Parametric Linear Programs

Consider

$$\begin{aligned} \max \quad & (3 \ 2 \ 0 \ 0) x \\ \text{s.t.} \quad & \begin{bmatrix} 3/2 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} 60 \\ 50 \end{bmatrix} \\ & x \geq 0 \end{aligned} \tag{P}$$

$B = \{1, 2\}$ is optimal basis. $\bar{x} = (20, 30, 0, 0)$. Then consider

$$\begin{aligned} \max \quad & (3 \ 2 \ 0 \ 0) x \\ \text{s.t.} \quad & \begin{bmatrix} 3/2 & 1 & 1 & 0 \\ 1 & 1 & 0 & 1 \end{bmatrix} x = \begin{bmatrix} 60 + \alpha \\ 50 - \alpha \end{bmatrix} \\ & x \geq 0 \end{aligned} \tag{P_\alpha}$$

For every α we get a different LP.

Define $g : \text{dom}(g) \rightarrow \mathbb{R}$ where

1. $\text{dom}(g) = \{\alpha : (P_\alpha) \text{ has an optimal solution}\}$.

2. $g(\alpha) = \text{OPT}(P_\alpha)$.

Goal is to describe g .

First we write (P_α) in canonical form for $B = \{1,2\}$

$$\begin{aligned} \max \quad & \bar{c}^T x + \bar{z} \\ \text{s.t.} \quad & \bar{A}x = \bar{b} \\ & x \geq 0 \end{aligned}$$

We then compute

$$\bar{A} = A_B^{-1}A = \begin{bmatrix} 1 & 0 & 2 & -2 \\ 0 & 1 & -2 & 3 \end{bmatrix}, \quad \bar{b} = A_B^{-1}b = \begin{bmatrix} 20 + 4\alpha \\ 30 - 5\alpha \end{bmatrix}$$

and

$$y = A_B^{-T}c_B = \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \quad \bar{c} = \begin{bmatrix} 0 \\ 0 \\ -2 \\ 0 \end{bmatrix}$$

$$\bar{z} = b^T y = 120 + 2\alpha$$

Then the tableau

$$T_{\{1,2\}} = \left[\begin{array}{cccc|c} 1 & 0 & 0 & 2 & 0 & 120 + 2\alpha \\ 0 & 1 & 0 & 2 & -2 & 20 + 4\alpha \\ 0 & 0 & 1 & -2 & 3 & 30 - 5\alpha \end{array} \right]$$

For what α is $B = \{1,2\}$ optimal? Equivalently, for what α is $B = \{1,2\}$ dual and primal feasible? For all values of α , it is dual feasible. We need to check for primal feasibility:

$$20 + 4\alpha \geq 0 \implies \alpha \geq -5$$

$$30 - 5\alpha \geq 0 \implies \alpha \leq 6$$

B optimal for all $\alpha \in [-5, 6]$.

Case $\alpha > 6$ Pivot (2,3) from $T_{\{1,2\}}$:

$$T_{\{1,3\}} \left[\begin{array}{cccc|c} 1 & 0 & 1 & 0 & 3 & 150 - 3\alpha \\ 0 & 1 & 1 & 0 & 1 & 50 - \alpha \\ 0 & 0 & -1/2 & 1 & -3/2 & -15 + 3\alpha/2 \end{array} \right]$$

For what α is $\{1,3\}$ optimal for (P_α) ?

$$50 - \alpha \geq 0 \implies \alpha \leq 50$$

$$-15 + \frac{5}{2}\alpha \geq 0 \implies \alpha \geq 6$$


So $\{1,3\}$ optimal if $\alpha \in [6, 50]$ and $g(\alpha) = 150 - 3\alpha$.

Case $\alpha > 50$ Try to run dual simplex, and notice primal is infeasible.

Case $\alpha < -5$ Similarly, we have

$$T_{\{2,4\}} \left[\begin{array}{cccc|c} 1 & 0 & 0 & 2 & 0 & 120 + 2\alpha \\ 0 & -1/2 & 0 & -1 & 1 & -10 - 2\alpha \\ 0 & 3/2 & 1 & 1 & 0 & 60 + \alpha \end{array} \right]$$

Then similarly, for $\alpha \in [-60, -5]$, $\{2,4\}$ is optimal. And $g(\alpha) = 120 + \alpha$.



PART III:

SOLVING OPTIMIZATION TECHNIQUES

Picture from Sdo, CC BY-SA 2.5 <https://creativecommons.org/licenses/by-sa/2.5>, via Wikimedia Commons

Cutting planes - review

relaxation

Consider optimization problems (P) & (Q) where

1. (P), (Q) both max/min & same objective function,
2. feasible region of (P) \supseteq feasible region of (Q)

Then (P) is a relaxation of (Q).

Key observation Suppose (P) is relaxation of (Q). If \bar{x} optimal for (P), \bar{x} feasible for (Q), then \bar{x} optimal for (Q).

Let (Q) be an IP. Let \bar{x} not feasible for (Q). Then $\alpha^T x \leq \beta$ is a cutting plane for \bar{x} if

1. $\alpha^T \bar{x} > \beta$ and
2. $\alpha^T x \leq \beta$ is valid for (Q).

“valid” here means for all feasible solution to (Q), $\alpha^T x \leq \beta$ holds.

10.1 Cutting plane algorithm

Consider

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \\ & x \text{ integer} \end{array} \quad (\text{Q})$$

and its LP relaxation

$$\begin{array}{ll} \max & c^T x \\ \text{s.t.} & Ax \leq b \end{array} \quad (\text{P})$$

Cutting plane algorithm:

1. If (P) infeasible, then (Q) infeasible. STOP
2. If (P) unbounded, then (Q) infeasible or unbounded (☹). STOP.
3. Find optimal solution \bar{x} of (P)

4. If \bar{x} integer, then \bar{x} optimal for (Q). STOP
5. adding cutting plane $\alpha^T x \leq \beta$ to $Ax \leq b$. Then repeat.

Denote LP relaxation by (P), (P)'s dual (D), LP after adding cutting plane by (P'), (P')'s dual (D'). Then the procedure is as follows: Solve (P), and get solution \bar{y} for (D). Then (\bar{y}) is feasible for (D'). Start from that feasible solution to get an optimal solution to (D'), then we have optimal solution for (P').

We know want to prove (☺). Suppose A, b, c rational.

Proposition 10.1

If (P) is unbounded, then (Q) is either infeasible or unbounded.

Lemma 10.2

$S = \{x : Mx \leq d\}$ where M and d are rational. If $S \neq \emptyset$, then S contains a rational point.

Proof of the proposition:

Suppose (P) unbounded, then there exists \bar{x}, r such that

1. $A\bar{x} \leq b$
2. $Ar \leq 0, c^T r > 0$

Let $S = \{r \mid Ar \leq 0, c^T r \geq 0\}$, by lemma, $\exists r' \in S$ which is integer. We may assume (Q) has a feasible solution x' . Then $x' + \lambda r'$ for $\lambda \in \mathbb{R}_+$ is feasible for (Q).

$$c^T(x' + \lambda r') = c^T x' + \lambda \underbrace{c^T r'}_{\geq 1} \rightarrow \infty$$

as $\lambda \rightarrow \infty$. □

Column generation

Goal is to solve

$$\begin{aligned} \min \quad & 1^T x \\ \text{s.t.} \quad & Mx \geq b, x \geq 0 \end{aligned} \quad (\text{P})$$

where M is $m \times n$ matrix, and n is huge.

Remark:

If (P) has an optimal solution, then it has an optimal solution with $\leq m$ non-zero variables.

Proof:

If (P) has an optimal solution, it has one that is an extreme point \bar{x} . \bar{x} is unique solution to set of tight constraints for \bar{x} . Then at least n tight constraints, at most m arise from $Mx \geq 0$. Then at least $n - m$ arise from $x \geq 0$. Then at most m non-zero entries for \bar{x} . \square

Restriction $J \subseteq \{1, \dots, n\}$.

$$\begin{aligned} \min \quad & 1^T s \\ \text{s.t.} \quad & M_J s \geq b, s \geq 0 \end{aligned} \quad (\text{P}_J)$$

Remark:

Let \bar{s} feasible for (P_J). Then \bar{x} is feasible for (P) where

$$\bar{x}_j = \begin{cases} \bar{s}_j & \text{if } j \in J \\ 0 & \text{otherwise} \end{cases}$$

11.1 Column generation algorithm

First attempt

1. Pick $J \subseteq \{1, \dots, n\}$.
2. Find an optimal solution \bar{s} for (P_J)
3. $\bar{x}_j = \begin{cases} \bar{s}_j & \text{if } j \in J \\ 0 & \text{otherwise} \end{cases}$
4. If \bar{x} optimal for (P), then STOP
5. pick j and set $J := J \cup \{j\}$. Then goto (2).

Lemma 11.1: Key lemma for column generation

Let \bar{s}, \bar{y} optimal for (P_J) and (D_J) . Define

$$\bar{x}_j = \begin{cases} \bar{s}_j & \text{if } j \in J \\ 0 & \text{otherwise} \end{cases}$$

If \bar{y} is feasible for (D) , then \bar{x} is optimal for (P) .

Proof:

We have \bar{x}, \bar{y} feasible for (P) and (D) . By weak duality, check whether $1^T \bar{x} \stackrel{?}{=} b^T \bar{y}$:

$$1^T \bar{x} = \sum_{j \in J} \bar{x}_j + \sum_{j \notin J} \bar{x}_j = 1^T \bar{s} = b^T \bar{y}$$

where the last equality is by strong duality: \bar{s}, \bar{y} optimal for (P_J) and (D_J) . □

Auxiliary problem

Input: \bar{y}

Output:

1. certificates \bar{y} feasible for (D) or
 2. find j such that the j^{th} constraint of (D) is violated for \bar{y} .
-

Recall the algorithm (slight different than last time):

1. Pick $J \subseteq \{1, \dots, n\}$.
2. \bar{s}, \bar{y} optimal for (P_J) and (D_J) .
3. $\bar{x}_j = \begin{cases} \bar{s}_j & \text{if } j \in J \\ 0 & \text{otherwise} \end{cases}$
4. If \bar{y} feasible for (D) , then STOP (\bar{x} optimal for (P))
5. Pic $\bar{j} \notin J$, such that constraint \bar{j} of (D) is violated for \bar{y} . $J := J \cup \{\bar{j}\}$.
6. Goto (2)

Step 4 and 5 is “auxiliary problem”: we define a function g whose domain is $\{1, \dots, n\}$, and $g(j) := \text{col}_j(M)^T \bar{y}$. We want to find \bar{j} maximize $g(j)$ over $j \in [n]$. If $g(\bar{j}) \leq 1$, then \bar{y} feasible for (D) . If $g(\bar{j}) > 1$, then constraint j of (D) is violated for \bar{y} .

11.2 Cutting stock

Given

- length of stock = 17m
- demand: $25 \times 3\text{m}$, $20 \times 5\text{m}$, $15 \times 9\text{m}$

We can have these cuts:

- (1) $3 + 3 + 3 + 3 + 3 + \dots$
- (2) $5 + 5 + 5 + \dots$

$$(3) 5 + 9 + \dots$$

$$(4) 9 + \dots$$

Then $5x(1)$, $2x(2)$, $14x(3)$, $1x(4)$, which satisfies the command. We used 22 stocks. This is not optimal. The goal is to meet demand and minimize the number of stocks. Then we can formulate as an IP.

There is an IP formulation with small number of variables and constraints. We cannot solve it well.

pattern for a stock

“Way of cutting stock into parts of length 3, 5 and 9”. Formally, the pattern is (u, v, w) where

u = number of parts of length 3

v = number of parts of length 5

w = number of parts of length 9

So the pattern of previous cuts are $(5, 0, 0)$, $(0, 3, 0)$, $(0, 1, 1)$, $(0, 0, 1)$. All possible patterns satisfy

$$3u + 5v + 9w \leq 17$$

where $u, v, w \geq 0$, integer.

Now let's write the IP formulation. Patterns

$$p_1 = \begin{pmatrix} u_1 \\ v_1 \\ w_1 \end{pmatrix}, \dots, p_n = \begin{pmatrix} u_n \\ v_n \\ w_n \end{pmatrix}$$

Then the optimization problem

$$\begin{aligned} \min \quad & \sum_{j=1}^n x_j \\ \text{s.t.} \quad & Mx \geq b = \begin{pmatrix} 25 \\ 20 \\ 15 \end{pmatrix} \\ & x \geq 0, x \text{ integer} \end{aligned}$$

where $M = [p_1 \mid p_2 \mid \dots \mid p_n]$. And denote its LP relaxation by (P).

Now let's do column generation on toy example. Let

$$p_1 = \begin{pmatrix} 5 \\ 0 \\ 0 \end{pmatrix}, \quad p_2 = \begin{pmatrix} 0 \\ 3 \\ 0 \end{pmatrix}, \quad p_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \quad J = \{1, 2, 3\}$$

We write (P_J) as before:

$$\begin{aligned} \min \quad & 1^T s \\ \text{s.t.} \quad & \begin{pmatrix} 5 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 1 \end{pmatrix} s \geq \begin{pmatrix} 25 \\ 20 \\ 15 \end{pmatrix} \\ & s \geq 0 \end{aligned} \tag{P_J}$$

Then we found $\bar{s} = (5, 20/3, 15)^T$, $\bar{y} = (1/5, 1/3, 1)^T$. Auxiliary problems:

$$g : \{\text{patterns}\} \rightarrow \bar{y}^T \begin{pmatrix} u \\ v \\ w \end{pmatrix}$$

Then we have new optimization problem

$$\begin{aligned} \max \quad & \frac{1}{5}u + \frac{1}{3}v + w \\ \text{s.t.} \quad & 3u + 5v + 9w \leq 17 \\ & u, v, w \geq 0, \text{ integer} \end{aligned}$$

This only has one constraint, which is knapsack problem. It can be solved efficiently. The optimal solution is $(1, 1, 1)^T$. Objective value > 1 . Then we add a new pattern $p_4 = (1, 1, 1)^T$ and $J = \{1, 2, 3, 4\}$ and solve again.